

Acid phosphatase observed in the intramembranous system of the yeast-like phase of *Paracoccidioides brasiliensis*

I. Campo-Aasen¹ & L. Yarzabal²

¹ Departments of Histochemistry and ² Immunoparasitology, Instituto Nacional de Dermatología, Facultad de Medicina, Universidad Central de Venezuela, Apartado 4043, Caracas 1010, Venezuela

In the present study the principles of the Gomori technique (3) combining cytochemical and electron microscopical methods according to the procedure of Holt & Hicks (4), were applied to the yeast form of *Paracoccidioides brasiliensis* (Pb). The fixative and substrate were changed according to the methodology of Etherton and Botham (2) and following the procedure of Borgers (1) for dealing with yeast. The motive was to observe the secretory process of the acid phosphatase as an indicator of the intrinsic protein secretion, and in consequence, to attempt the demonstration of the endoplasmic reticulum and the Golgi apparatus of these cells.

The acid phosphatase spherical deposits are observed mostly in the matrical side of the plasmalemma. They seem to be more intense at 5 days culture. They are varied in size and opacity. The general structure of the fungi is preserved. The controls were negative and no activity was observed in the inhibition with sodium fluoride and tartatic acid 0.01 M.

We do not consider that we have demonstrated the endoplasmic-reticulum and the Golgi apparatus with the acid phosphatase technique, but we are prone to think instead that the intramembranous structures of the fungi derived from the internal phase of its plasma membrane are rich in one of the multiple forms or isoenzymes of the acid phosphatase (5).

DDCT-SAL-3 of Consejo Nacional de Investigaciones Científicas y Tecnológicas (CONICIT), Venezuela.

We gratefully acknowledge the technical assistance of Miss Z. Cova and N. Ghersy.



Fig. 1. *P. brasiliensis*. 5 days culture, yeast form. Spherical deposits of acid phosphatase of different sizes and densities are observed in the plasma membrane (PM) and in intramembranous structures (IS). 40 000X.

Acknowledgements

This work has been supported by Grant N°




Fig. 2. *P. brasiliensis*. 5 days culture, yeast form. 2a. Intramembranous structures (IS) with deposits of acid phosphatase at 70 000X. 2b. Intramembranous structures (IS) enlarged to 100 000X. The acid phosphatase are observed mostly at the matrical side of the membranes (arrows).

References

1. Borgers, M. & de Nollin, S., 1974. The preservation of subcellular organelles of *Candida albicans* with conventional fixatives. *J. Cell. Biol.* 62: 574-581.
2. Etherton, J. E. & Botham, C. M., 1970. Factors affecting lead capture methods for the fine localization of rat lung acid phosphatase. *Histochem. J.* 2: 507-519.
3. Gomori, G. *Microscopic Histochemistry: Principles and Practice*. p. 189. Chicago University Press, Chicago 1952.
4. Holt, S. J. & Hicks, R. M., 1961. The localization of acid phosphatase in rat liver cells as revealed by combined cytochemical staining and electron microscopy. *J. Biophys. Biochem. Cytol.* 11: 47-66.
5. Markert, C. L. & Apella, E., 1961. In: *Multiple molecular forms of enzymes*. Ed. F. Wroblewski, Ann. New York Acad. Sci. 94: 678-689.